

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



File

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/68, C12N 15/00, C07H 21/04</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/39536</b> <b>(43) International Publication Date:</b> 12 December 1996 (12.12.96)
<b>(21) International Application Number:</b> PCT/US96/08142 <b>(22) International Filing Date:</b> 31 May 1996 (31.05.96) <b>(30) Priority Data:</b> 08/467,909 6 June 1995 (06.06.95) US <b>(71) Applicant (for all designated States except US):</b> T CELL DIAGNOSTICS, INC. [US/US]; 115 Fourth Avenue, Needham, MA 02194 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> EMMETT, Constance [US/US]; 18 Pratt Street, Melrose, MA 02176-1930 (US). FOSTER, Kimberly, A. [US/US]; 641 Main Street, Boylston, MA 01505 (US). <b>(74) Agents:</b> LICATA, Jane, Massey et al.; Suite 201, 210 Lake Drive East, Cherry Hill, NJ 08002 (US).			<b>(81) Designated States:</b> AL, AM, AU, BB, BG, BR, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> With international search report.
<b>(54) Title:</b> UNIVERSAL CHEMISTRY ENZYME-LINKED IMMUNOSORBENT ASSAY FOR DETECTION OF mRNA EXPRESSION			
<b>(57) Abstract</b> <p>A kit for the quantitation of mRNA expression levels in a biological sample is provided which contains a detergent-chaotrope lysis mixture for extracting total RNA from the biological sample; a mixture of reverse transcriptase, nucleotides and an oligo-dT primer to synthesize cDNA from polyadenylated mRNA in the total mRNA; a means for labeling a primer or probe with a molecule which binds to an agent coated on a microtiter plate; a microtiter plate coated with an agent capable of binding to said molecule; a means for labeling a primer or probe with a detectable label; an antibody which detects the labeled primer or probe; and a means for detecting the antibody. Methods for quantitating mRNA expression in a biological sample are also provided.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**UNIVERSAL CHEMISTRY ENZYME-LINKED IMMUNOSORBENT ASSAY  
FOR DETECTION OF mRNA EXPRESSION**

**BACKGROUND OF THE INVENTION**

Polymerase chain reaction (PCR) has been recognized as  
5 a powerful technology for the characterization and quantitation  
of DNA. Recent methodologies for the direct performance of PCR  
on washed blood cells and whole blood significantly increase  
its usefulness for the processing and characterization of large  
numbers of blood samples. Wu et al. *Transgenica* 1994, 1(1):35-  
10 38.

Analyses of PCR products using an enzyme-linked  
immunosorbent assay have been disclosed by several  
investigators. For example, Rasmussen et al., *Clinical  
Chemistry* 1994, 40(2):200-205 teach detection of an  
15 immobilized, amplified product in a one phase system. This  
assay integrates the polymerase chain reaction with  
hybridization of the amplified product for detection in the  
same microwell. Detection of bovine leukemia virus and  
Salmonella was demonstrated, however, it is suggested that the  
20 assay is easily adaptable for other organisms, simply by using  
other primers and probes. However, because of poor  
reproducibility, this assay is recommended only for qualitative  
analysis. Genemed Biotechnologies, Inc. (South San Francisco,  
CA) also provides several ColorTect® enzyme-based signal  
25 generation kits which can be used to study several PCR samples  
simultaneously using a 96 microwell plate format.

The quantitation of RNA can also be performed by a  
similar technique, however, using RNA as the starting material

- 2 -

requires a reverse transcription step to synthesize a cDNA template for PCR. This method is referred to as RT-PCR. Quantitation of messenger RNA is achieved by RT-PCR.

A reverse transcription (RT) *in situ* PCR protocol in which direct incorporation of digoxigenin-11-dUTP allows the detection of PCR amplified RNA (via cDNA) in intact cells is described by Nuovo, G.J., *Biochemica* 1994, 11(1):4-6. In this method, after digesting all non-target DNAs with RNase-free DNase, the desired cDNA is specifically amplified by first performing reverse transcription with a downstream primer and then *in situ* PCR with digoxigenin-11-dUTP. Incorporated digoxigenin-11-dUTP is then detected with alkaline phosphatase-conjugated antidigoxigenin and the colorimetric alkaline phosphatase substrates NBT and BCIP. This method of RT *in situ* PCR was compared to standard *in situ* hybridization and found to be more sensitive in the detection of hepatitis C virus. It is suggested that the detection of digoxigenin-labeled PCR-amplified viral RNAs and mRNAs can provide information not obtainable with either solution-phase PCR or standard *in situ* hybridization.

It has now been found that mRNA expression levels for various proteins can be quantitated in biological samples using a single assay which combines RT-PCR with ELISA. The determination of mRNA expression levels can be used in the early diagnosis of disease states and in monitoring the efficacy of potential therapeutics for these diseases.

#### SUMMARY OF THE INVENTION

An object of the present invention is to provide a kit for the quantitation of mRNA expression levels in biological samples comprising a detergent-chaotrope lysis mixture for extracting total RNA from a biological sample; a mixture of reverse transcriptase, nucleotides and an oligo-dT primer to synthesize cDNA from the polyadenylated mRNA in the total mRNA population; a means for labeling a primer or probe with a molecule which binds to an agent coated on a microtiter plate; a means for labeling a primer or probe with a detectable label;

- 3 -

an antibody which detects the labeled primer or probe; a means for detecting the antibody; and a coated microtiter plate.

Another object of the present invention is to provide a method for quantitating levels of mRNA expression. This method  
5 comprises extracting total RNA from a biological sample. cDNA is then synthesized from mRNA in the extracted total RNA. Nucleotides, Taq polymerase, a magnesium source and a 5' and 3' primer selected in accordance with the mRNA to be quantitated, wherein the 5' primer is labeled with either a molecule capable  
10 of binding to a coated microtiter plate or a detectable marker, are then added to the cDNA and a PCR reaction is performed to obtain a PCR product. A probe is then added to the PCR product under conditions in which the probe hybridizes to the PCR product, said probe being capable of hybridizing to the PCR  
15 product and containing either (1) a molecule capable of binding to a coated microtiter plate, or (2) a detectable marker. The sample which contains a mixture of hybridized probe and PCR product, unhybridized probe and PCR products, and excess primers is then transferred to a microtiter plate coated with  
20 an agent capable of binding to the molecule on the PCR product or probe under conditions in which this molecule will bind to the microtiter plate. Detectable marker on bound hybridized probe and PCR product is then detected and mRNA levels determined.

## 25 DETAILED DESCRIPTION OF THE INVENTION

In many disease states it is faster, as well as more revealing, to track the mRNA expression levels of a selected protein for diagnosis of the disease as well as monitoring of therapies for the disease rather than wait for the production  
30 of the selected protein. In other cases, such as transplantation rejection, the event has occurred before the diagnosis can be made and a cytokine protein can be detected. The detection of expression levels of cytokine mRNA can precede this and be revealed while the rejection process can still be  
35 mitigated.

- 4 -

In the present invention, an assay is provided for the detection of mRNA expression of a selected protein in a biological sample using a universal enzyme-linked immunosorbent assay (ELISA) or probed polymerase chain reaction (PCR) products resulting from reverse transcriptase-PCR (RT-PCR) of the mRNA of interest. By "biological sample" it is meant to include, but is not limited to, whole blood, serum, plasma, urine, synovial fluid, saliva, cerebrospinal fluid, tissue infiltrate, cervical or vaginal exudate, pleural effusion, bronchioalveolar lavage fluid, gastric lavage fluid, small or large bowel contents, and swab specimens from various bodily orifices dispersed in a suitable medium. In a preferred embodiment, whole blood is used. In a preferred embodiment, detection of the mRNA is performed by well known colorimetric or fluorometric methods. The assay is constructed in a universal chemistry format wherein different RT-PCR primers and probes can be used for the detection of mRNA of a selected protein.

For example, in one embodiment, the present invention can be used to detect cellular levels of cytokine mRNA such as interleukin-8 (IL-8) messenger RNA in whole blood. PCR primers for cytokines such as  $\beta$ -actin, IL-1 $\alpha$ , IL-1 $\beta$ , TNF- $\alpha$ , IL-6, IL-8 and GM-CSF which can be used in the present invention are described by Zhong et al., *Arch. Surg.* 1993, 128:158-163.

In another embodiment, the present invention can be used to detect the cytokines, interleukin-2 (IL-2) and interferon- $\gamma$  (IFN- $\gamma$ ). Elevated mRNA expression levels of these cytokines can be used to detect acute rejection of renal transplants prior to any other present method of diagnosis. Lipmann et al., *Transplantation* 1992, 53:73-79. Primer and probe sequences which can be used for the detection of IL-2 include: the 5' primer 5'-CATTGCACTAAGTCTTGCACTTGTC-3' (SEQ ID NO: 1); the 3' primer 5'-CGTTGATATTGCTGATTAAGTCCCTG-3' (SEQ ID NO: 2); and the probe 5'-TTCTTCTAGACACTGAAGATGTTTCAGTTC-3' (SEQ ID NO: 3). Primer and probe sequences which can be used for the detection of IFN- $\gamma$  include: the 5' primer 5'-GCATCGTTTTGGGTTCTCTTGGCTGTTACTGC-3' (SEQ ID NO: 4); the 3'

- 5 -

primer 5'-CTCCTTTTTCGCTTCCCTGTTTTAGCTGCTGG-3' (SEQ ID NO: 5);  
and the probe 5'-TCTGGTCATCTTTAAAGTTTTTAAAAAGTT-3' (SEQ ID NO:  
6). In a preferred embodiment, the 5' primer is labeled with  
biotin at the 5' end of the sequence and the probe is labeled  
5 with fluorescein isothiocyanate (FITC) at the 5' end of the  
sequence.

In yet another embodiment, the present invention can be  
used to detect placental isoform of ferritin (PLF) messenger RNA. PLF  
mRNA expression levels have been related to the development of  
10 breast cancer. Breast cancer is difficult to diagnose at an  
early stage, when most amenable to therapy, and knowledge about  
the expression levels of PLF mRNA is believed to be a  
definitive way to diagnose and begin therapy as soon as  
possible. The efficacy of the therapy can also be monitored by  
15 following levels of mRNA PLF using the present invention.

In an assay performed in accordance with the present  
invention, total RNA is extracted from a biological sample. A  
mixture comprising reverse transcriptase, which is RNA-  
dependent DNA polymerase, nucleotides and an oligo-dT primer is  
20 then added to the extracted RNA to synthesize cDNA from the  
polyadenylated mRNA in the total mRNA population. In the same  
tube in which the cDNA is synthesized, more nucleotides, Taq  
polymerase, which is thermally stable DNA-dependent polymerase,  
a magnesium source and two primers selected in accordance with  
25 the mRNA to be detected are added and the PCR reaction run.  
The two primers are designed to prime inward from the ends of  
the known sequence of cDNA which represents the salient portion  
of the gene. In a first embodiment, the 5' primer is labeled  
with a molecule capable of binding to a coated microtiter  
30 plate. For example, the 5' primer can be labeled with biotin  
and the microtiter plate coated with streptavidin or the 5'  
primer can be labeled with digoxigenin and the microtiter plate  
coated with anti-digoxigenin antibody. In this embodiment, a  
probe designed to complement a selected portion of the target  
35 gene is then added to the tube. By "selected portion", it is  
meant that portion of the gene sequence which is not primed by  
the PCR primers. The probe is labeled with a detectable marker

- 6 -

such as FITC or digoxigenin and is capable of hybridizing to the labeled PCR product. In a second embodiment, the 5' primer is labeled with a detectable marker such as fluorescein isothiocyanate (FITC) or digoxigenin. In this embodiment, the probe, also designed to complement a selected portion of the target gene, is labeled with a molecule capable of binding to a coated microtiter plate. For example, the probe can be labeled with biotin and the microtiter plate coated with streptavidin or the probe can be labeled with digoxigenin and the microtiter plate coated with anti-digoxigenin antibody. The mixture of hybridized probe and PCR product, unhybridized probe and PCR products and excess primers are then added to a microtiter plate coated with an agent capable of binding to the molecule on either the PCR product or the probe. In a preferred embodiment, the microtiter plate is coated with streptavidin. A detectably labeled antibody capable of binding to any labeled PCR product which is hybridized to the probe and bound to the plate is then added and the antibody is detected. In a preferred embodiment, the antibody is detected colorimetrically or fluorometrically. For example, in one embodiment wherein the PCR product or probe is labeled with FITC, the antibody may comprise anti-FITC horse radish peroxidase which is detectable colorimetrically by the addition of 3,3',5,5'-tetramethylbenzidine (TMB) peroxidase substrate or O'-phenylenediamine (OPD). In another embodiment wherein the PCR product or probe is labeled with FITC, the antibody may comprise anti-FITC alkaline phosphatase which is detectable fluorometrically by the addition of methylumbelliferyl phosphate (MUP).

In the present invention kits comprising components used in this universal enzyme-linked immunosorbent assay of probed PCR products resulting from reverse transcriptase-PCR (RT-PCR) of a particular mRNA of interest are also provided. Kits of the present invention are constructed using universal chemistry format in that the only differences are the RT-PCR primers and probe for the targets of interest. The remainder of the assay and kit components remain the same and can be routinely used



- 7 -

with various primers and probes for any target of interest. The construction of these kits in a universal chemistry manner provides significant advantage for the research and clinical setting.

5       Kits of the present invention comprise a detergent-chaotrope lysis mixture for extracting total RNA from a biological sample; a mixture of reverse transcriptase, nucleotides and an oligo-dT primer to synthesize cDNA from the polyadenylated mRNA in the total mRNA population; a means for  
10 labeling a primer or probe with a molecule which binds to an agent coated on a microtiter plate; a means for labeling a primer or probe with a detectable label, an antibody which detects the labeled primer or probe; a means for detecting the antibody; and a microtiter plate coated with an agent capable  
15 of binding to the molecule on either the probe or the PCR product. With this kit, the investigator need only provide the specific primers and probes to be used for a selected target to be measured.

20       The following nonlimiting examples are provided to further illustrate the present invention

#### EXAMPLES

##### Example 1: Isolation of RNA from whole blood

Whole blood (50  $\mu$ l) was pelleted and washed twice with a NaCl-EDTA buffer in the following manner. The whole blood  
25 was mixed with 500  $\mu$ l of 10 mM NaCl-EDTA buffer, vortexed vigorously and spun in a microcentrifuge at 14,000 for 5 minutes. The supernatants were poured off and the pellets washed a second time in the NaCl-EDTA buffer. Following a second centrifugation, the pellets were resuspended in 500  $\mu$ l  
30 denaturation solution (5 M guanidinium thiocyanate, 0.125 M Tris-HCl (pH 7.4), 0.3125 M sodium acetate, and 1.25%  $\beta$ -mercaptoethanol), 500  $\mu$ L TRAx<sup>®</sup> lysis buffer (1.56 dibasic potassium phosphate, 0.14 g monobasic potassium phosphate, 8.77 g sodium chloride, 90 ml Triton-X and 60 ml Nonidet P-40) and  
35 100  $\mu$ g tRNA. The lysis mixture was vigorously vortexed and the tubes incubated at 65°C for 10 minutes. The tubes were iced

- 8 -

for at least 1 minute, vortexed, and then split into a fresh tube so that 500  $\mu$ l isopropanol could be added to each tube. The tubes were vortexed at high speed for 10 seconds each, then centrifuged at 14,000 RPM for 15 minutes. The supernatants  
5 were poured off, and the visible white pellets washed with 1.5 ml 70% ethanol and centrifuged at 14,000 RPM for 10 minutes. The supernatants were poured off and the pellets dried in a Speed-Vac with no heat for one hour. The dried pellets were resuspended in 50  $\mu$ l sterile water and frozen at -70°C. Before  
10 use, the pellets were heated at 50°C-65°C until they dissolved in water.

**Example 2: Synthesis of cDNA**

A mixture of reverse transcriptase, which is RNA-dependent DNA polymerase, nucleotides and an oligo-DT primer  
15 are added to synthesize cDNA from the polyadenylated mRNA in the isolated total RNA population. This procedure requires temperature which are achieved in the same thermal cycler as described in Example 3.

**Example 3: PCR reaction**

20 In the same tube in which the cDNA is synthesized, additional nucleotides, Taq polymerase, a source of magnesium and the two primers are added and PCR reaction run in the thermal cycler for 35 cycles. The primers are designed to prime inwardly from the ends of the known sequence of cDNA  
25 which represents the salient part of the gene. The 5' primer is biotin-labeled and the 3' primer is unlabelled. The resultant biotin-labelled PCR products are hybridized to the fluorescein isothiocyanate (FITC)-labelled probe in the same tube. The probe is designed to complement a selected portion  
30 of the target sequence which is not primed by the PCR primers.

This PCR reaction is asymmetrical yielding a single-stranded PCR product. This reaction is accomplished by using an excess amount of biotin-labelled 5' primer (50  $\mu$ M) and a limiting amount of the unlabelled 3' primer (1  $\mu$ M). The

- 9 -

resultant single-stranded, biotin-labelled PCR product then easily hybridizes with a FITC labeled probe.

**Example 4: ELISA assay**

The mixture of hybridized FITC-labelled probe and biotin-  
5 labelled PCR product, unhybridized probes and PCR products and  
excess primers are added to a microtiter plate coated with  
streptavidin. All versions of the biotin-labelled primer or  
PCR product bind to the streptavidin quickly and very strongly  
during a 30 minute room temperature incubation. Subsequent  
10 wash steps remove all unhybridized probes and unlabeled  
primers. The FITC-labeled probes hybridize only to the full  
length biotin-labeled PCR products, and only those which are  
hybridized to the bound-labelled PCR products remain in the  
wells after the washes.

15 The bound hybrid, consisting of FITC-labelled probe and  
the biotin-labelled PCR product, bind sheep anti-FITC  
horseradish peroxidase (HRP) conjugated during incubation at  
room temperature. After another wash step, the TMB-peroxidase  
substrate (3,3', 5,5'-tetramethylbenzidine) is added and the  
20 color is allowed to develop. The reaction is stopped by the  
addition of a weak acid solution. The plate is then read in an  
ELISA plate reader.

**Example 5: Determining IL-8 mRNA Expression**

mRNA was isolated from whole blood as described in  
25 Example 1. The RNA pellet was resuspended in 20  $\mu$ l of RNase-  
free water. The RNA suspension (10  $\mu$ l) plus oligo dT primer (1  
 $\mu$ l; 0.5  $\mu$ g/ $\mu$ l) and RNase-free water (2  $\mu$ l) were incubated at  
70°C for 10 minutes. The samples were then placed on ice for  
1 minute and 7  $\mu$ l of master mix containing 10X PCR buffer, 25  
30 mM MgCl<sub>2</sub>, 10 mM each dNTP mix and 0.1 M DTT was added. The  
samples were incubated at 42°C for 5 minutes. Reverse  
transcriptase (1  $\mu$ l) was added and the samples were incubated  
again at 42°C for 50 minutes, followed by a 15 minute  
incubation at 70°C. The samples were then chilled on ice.

- 10 -

The asymmetric PCR reaction was then performed using 10X PCR buffer, 25 mM MgCl<sub>2</sub>, 10 mM each dNTP, Taq polymerase (2 units/reaction), 50 μM of human IL-8 5' primer: 5'-biotin-ATGACTTCCAAGCTGGCCGTGGCT-3' (SEQ ID NO: 7), and 1 μM human IL-8 3' primer: 5'-TCTCAGCCCTCTTCAAAACTTCTC-3' (SEQ ID NO: 8) for 30 to 35 cycles.

Thirty microliters of the resulting PCR product were then hybridized with 20 μl of a hybridization mixture containing 100 mM HEPES, 0.1% Tween 20, 200 μM EDTA, 1 M NaCl and 20 pmol of a human IL-8 probe: 5'-FITC-CACAGAGCTGCAGAAATCAGGAAGGCTGCCA-3' (SEQ ID NO: 9). The mixture was then incubated at 95°C for 5 minutes followed by a 1.5 hour incubation at 37°C. An ELISA assay was performed in the hybridized mixture as described in Example 4.

- 11 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(i) APPLICANT: Constance Emmett, Kimberly A. Foster

(ii) TITLE OF INVENTION: Universal Chemistry Enzyme-Linked  
Immunosorbent Assay for Detection of  
mRNA Expression

(iii) NUMBER OF SEQUENCES: 9

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Jane Massey Licata, Esq.

(B) STREET: 210 Lake Drive East, Suite 201

(C) CITY: Cherry Hill

(D) STATE: NJ

(E) COUNTRY: USA

(F) ZIP: 08002

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: DISKETTE, 3.5 INCH, 1.44 Mb STORAGE

(B) COMPUTER: IBM 486

(C) OPERATING SYSTEM: WINDOWS FOR WORKGROUPS

(D) SOFTWARE: WORDPERFECT 5.1

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: not yet assigned

(B) FILING DATE: Herewith

(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Jane Massey Licata

(B) REGISTRATION NUMBER: 32,257

(C) REFERENCE/DOCKET NUMBER: TCEL-0039

- 12 -

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (609) 779-2400

(B) TELEFAX: (609) 779-8488

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CATTGCACTA AGTCTTGAC TGTCA 26

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CGTTGATATT GCTGATTAAG TCCCTG 26

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

- 13 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

TTCTTCTAGA CACTGAAGAT GTTTCAGTTC 30

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GCATCGTTTT GGGTTCCTCTT GGCTGTTACT GC 32

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CTCCTTTTTTC GCTTCCCTGT TTTAGCTGCT GG 32

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

- 14 -

TCTGGTCATC TTAAAGTTT TAAAAAGTT

30

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

ATGACTTCCA AGCTGGCCGT GGCT

24

## (2) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TCTCAGCCCT CTTCAAAAAC TTCTC

25

## (2) INFORMATION FOR SEQ ID NO: 9:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CACAGAGCTG CAGAAATCAG GAAGGCTGCC A

31



- 15 -

**What is claimed is:**

1. A kit for the quantitation of mRNA expression levels in a biological sample comprising:
  - a) a detergent-chaotrope lysis mixture for  
5 extracting total RNA from a biological sample;
  - b) a mixture of reverse transcriptase, nucleotides and an oligo-dT primer to synthesize cDNA from polyadenylated mRNA in the total mRNA;
  - c) a means for labeling a primer or probe with a  
10 molecule which binds to an agent coated on a microtiter plate;
  - d) a microtiter plate coated with an agent capable of binding to said molecule;
  - e) a means for labeling a primer or probe with a detectable label;
  - 15 f) an antibody which detects the labeled primer or probe; and
  - g) a means for detecting the antibody.
2. A method for quantitating levels of mRNA expression  
20 comprising:
  - a) extracting total RNA from a biological sample;
  - b) synthesizing cDNA from mRNA in the extracted total RNA;
  - c) adding nucleotides, Taq polymerase, a magnesium  
25 source and a 5' and 3' primer selected in accordance with the mRNA to be detected to the cDNA, wherein said 5' primer is labeled with a molecule capable of binding to a coated microtiter plate;
  - d) performing a PCR reaction to obtain a PCR  
30 product;
  - e) adding a probe to the PCR product which is capable of hybridizing to the PCR product and labeled with a detectable marker under conditions in which the probe hybridizes to the PCR product;
  - 35 f) transferring a mixture of hybridized probe and PCR product, unhybridized probe and PCR products and excess primers to a microtiter plate coated with an agent capable of

- 16 -

binding to the molecule on the PCR product under conditions in which the molecule on the PCR product will bind to the microtiter plate; and

g) detecting the detectable marker on bound  
5 hybridized probe and PCR product so that mRNA levels can be determined.

3. A method for quantitating levels of mRNA expression comprising:

a) extracting total RNA from a biological sample;  
10 b) synthesizing cDNA from mRNA in the extracted total RNA;

c) adding nucleotides, Taq polymerase, a magnesium source and a 5' and 3' primer selected in accordance with the mRNA to be detected to the cDNA, wherein said 5' primer is  
15 labeled with a detectable marker;

d) performing a PCR reaction to obtain a PCR product;

e) adding to the PCR product a probe capable of hybridizing to the PCR product and containing a molecule  
20 capable of binding to an agent coated on a microtiter plate under conditions in which the probe will bind to the PCR product;

f) transferring a mixture of hybridized probe and PCR product, unhybridized probe and PCR products and excess  
25 primers to a microtiter plate coated with an agent capable of binding to the molecule on the probe under conditions in which the molecule on the probe will bind to the microtiter plate; and

g) detecting the detectable marker on bound  
30 hybridized probe and PCR product so that mRNA levels can be determined.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/08142

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; C12N 15/00; C07H 21/04  
US CL : 435/6, 172.3; 536/24.3

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 172.3, 810; 436/501; 514/44; 536/24.3; 935/77, 78

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MANIATIS et al., "Molecular Cloning, A laboratory manual". New York: Cold Spring Harbor Laboratory, 1982, page 212-216 and 280-281, see entire document.	1-3
Y	MATTHEWS et al. Review, Analytical Strategies for Use of DNA Probes. Analytical Biochemistry, 1988, Volume 169, pages 1-25, especially page 1 and Figure 10 page 17.	1-3
Y	SYVANEN et al. Fast quantification of nucleic acid hybrids by affinity-based hybrid collection. Nucleic Acids Research, 1986, Volume 14, Number 12, pages 5037-5048. See whole document.	1-3

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 JUNE 1996

Date of mailing of the international search report

12 JUL 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEZIA RILEY

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/08142

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SYVANEN et al. Quantification of polymerase chain reaction products by affinity-based hybrid collection. Nucleic Acids Research, 1988, Volume 16, Number 23, pages 11327-11338. See whole document.	1-3

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/08142

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, STN

search terms: Nucleic, PCR or Polymerase chain reaction, quantitative, matrix, solid support